



PRODUCT DATA SHEET

LAV-CIDE

Lav-cide is designed specifically as a general cleaner and disinfectant for use on hard, non-porous surfaces in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, veterinary clinics, pet shops, animal life science laboratories and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food. In addition, Lav-cide deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

Lav-cide is effective against the virus that causes COVID-19 on hard, non-porous surfaces when used according to disinfection directions

DIRECTIONS FOR USE:

Disinfecting, deodorizing and cleaning: Remove visible soil prior to application of the product. Hold container six to eight inches from surface to be treated. Allow product to wet the hard non-porous surface for 3 minutes. No scrubbing is necessary. Wipe with a clean cloth or allow to air dry. Lav-cide will not leave grit or soap scum.

Bathroom Disinfectant Cleaner: Lav-cide is also a bowl and bathroom cleaner that cleans, disinfects and deodorizes. Cleans, disinfects and deodorizes sinks, sink rims, sink basins, faucets, tubs, glazed tiles, glazed ceramic, glazed porcelain, chrome, stainless steel, and all non-porous inanimate surfaces found in the bathroom. Eliminates odors, leaving restrooms smelling clean and fresh. Cleans, disinfects, and deodorizes hard non-porous surfaces by killing odor-causing microorganisms and controls the growth of non-pathogenic mold and mildew for up to 7 days. Use daily on: glass, glazed ceramic, glazed tile, glazed porcelain, chrome, stainless steel, and plastic surfaces associated with floors, walls, fixtures, toilets and urinals when used externally or above waterline only, sinks, shower rooms, and locker rooms.

EPA Number 1839-83-1674

See reverse side for efficacy data.

To disinfect food service establishment food contact surfaces: Before using, food products and packaging materials must be removed or carefully protected. For countertops, appliances, tables, hold container six to eight inches from the surface to be treated. Allow Lav-cide to wet the hard non-porous surface for 3 minutes. Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Visibly soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

SPECIFICATIONS

Appearance	Blue transparent liquid
Fragrance added	Floral
pH	12-12.5
Specific gravity @25°	0.90-1.08
Unopened shelf life	1 year

Refer to the Safety Data Sheet (SDS) for detailed safety, health and environmental information.

PACKAGING

Available in 1- quart plastic bottles with squirt top (12/case).

January 2024



11 Norfolk Street
Mansfield, MA 02048
877-4-JANSAN
www.nextgensupply.com

Next-Gen Supply Group is part of Envoy Solutions company, a specialized distributor and solution provider in jan-san, foodservice, packaging, and marketing execution.

BACTERICIDAL ACTIVITY: Using AOAC test methods in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, Lav-cide kills the following organisms on hard non-porous inanimate surfaces:

Organism	Contact Time
Pseudomonas aeruginosa	3 minutes
Staphylococcus aureus	3 minutes
Salmonella (choleraesuis) enterica	3 minutes
Escherichia coli	3 minutes
Escherichia coli O157:H7	3 minutes
Streptococcus pyogenes (V.A. Medical Center Isolate 04001)	3 minutes
Listeria monocytogenes	3 minutes
Yersinia enterocolitica	3 minutes
Enterococcus faecium	3 minutes
Corynebacterium ammoniagenes	3 minutes
Salmonella (typhi) enterica	3 minutes
Shigella flexneri	3 minutes
Corynebacterium bovis	3 minutes
Shigella dysenteriae	3 minutes

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: Using AOAC test methods in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, Lav-cide kills the following organisms on hard non-porous inanimate surfaces:

Organism	Contact Time
Methicillin resistant Staphylococcus aureus (MRSA)	3 minutes
Vancomycin resistant Enterococcus faecalis (VRE)	3 minutes
Vancomycin intermediate resistant Staphylococcus aureus (VISA)	3 minutes
Methicillin resistant Staphylococcus epidermidis (MRSE)	3 minutes
Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA)	3 minutes
Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA)	3 minutes

+VIRUCIDAL ACTIVITY (Viruses)

Virus Enveloped

	Contact Time
HIV-1 (associated with AIDS)	1 minute
Avian Influenza A Virus strain H3N2	2 minutes
Avian Influenza A Virus strain H9N2	2 minutes
Hepatitis B Virus (HBV)	1 or 5 minutes
Hepatitis C Virus (HCV)	1 or 5 minutes
Duck Hepatitis B Virus (DHBV)	1 or 5 minutes
Bovine Viral Diarrhea Virus (BVDV)	1 or 5 minutes
Human Coronavirus	2 minutes
SARS associated Coronavirus	2 minutes
SARS-Related Coronavirus 2 [SARS-CoV-2] USA-WA1/2020	1 minute
Rabies Virus	30 seconds
Paramyxovirus (Mumps)	3 minutes
Influenza A Virus H1N1	1 minute

Virus Large Non-Enveloped

Rotavirus	3 minutes
-----------	-----------

Virus Small Non-Enveloped

Norovirus (Norwalk virus),	30 seconds
Feline Calicivirus	30 seconds
Canine Parvovirus	10 minutes
Poliovirus Type 1	10 minutes
Rhinovirus type 39	3 minutes
Hepatitis A Virus (HAV) (University of Ottawa)	10 minutes

Fungi

	Contact Time
Trichophyton interdigitale, formerly Trichophyton mentagrophytes	10 minutes